

Open

Farmacos betalactamicos pdf

Situación	Agente	Régimen
régimen estándar de profilaxis	**Amoxicilina**	Adultos: 2 g niños: 50 mg/kg oral 1 hora antes del procedimiento oral.
Si no se puede vía oral	**Ampicilina**	Adultos: 2g IM o IV niños: 50mg/kg IM, IV 30 minutos antes del procedimiento oral.
Alérgicos a la penicilina	**Clindamicina**	Adultos:600mg Niños: 20 mg/kg oral 1 hora antes del procedimiento oral.
	cephalexin o cefadroxil	Adultos: 2 g niños: 50mg/kg oral 1 hora antes del procedimiento oral.
	Clindamicina	adultos: 600 mg Niños: 20mg/kg IV en 30 minutos antes del procedimiento oral.
alergia a la penicilina e imposibilidad de usar vía oral	**Clindamicina**	adultos: 600 mg Niños: 20mg/kg IV en 30 minutos antes del procedimiento oral.
	Cefazolin	Adultos:1.0g Niños: 25 mg/kg IM o IV En 30 minutos antes del procedimiento oral.



Tabla V. Betalactámicos que comparten cadenas laterales

Penicilinas	*Cefalosporinas*	
		Generación
Amoxicilina	Cefadroxilo	1ª
	Cefprozil	
	Cefalexina	
Ampicilina	Cefalexina	1ª
	Cefadrina	
	Loracarbef	2ª
	Cefaclor	
Piperacilina	Cefoperazona	3ª
Bencilpenicilina	Cefalotina	1ª
	Cefamandol	
	Cefaloridina	
CEFALOSPORINAS		
Ceftriaxona (3ª)	Cefotaxima	3ª
	Cefpodoxima	
	Cefditoren	
	Ceftazidina	
Cefoxitina (2ª)	Cefalotina	1ª
	Cefuroxima	2ª
Cefamandol (2ª)	Cefonicid	2ª
	Cefotetam	
MONOBACTAM		
Aztreonam	Ceftazidima	3ª

Lista de farmacos betalactamicos. Farmacos betalactamicos ejemplos. Grupo de farmacos betalactamicos. Farmacos antibioticos betalactamicos. Farmacos betalactamicos pdf.

alla atattada e ocinilc etnemlatnemadnof otnematneiro id ,icittalateb ied emeisniâlled elauttecnoc enoisiv anu erirffo id olleuq Ì Ã olocitra otseuq id ovitteiboâL. Ãtilibisnes attodir noc imsinagrorcim ad etasuac inoizefni enucla id otnemattart len icitoibitna itseuq id icimanidocamraf e icitenicocamraf irtemarap i erazzimitto da arim ehc acitueparet aigetarts anu id etrap emoc aunitnoc enoisufni ni icimttal-ateb id enoizartsinimmos al rep esseretni etnecserc li Ì Ã inna imitlu ilged Ãtivon ednarg al esroF .giF(.attircsed arpos arutturts al eramrof rep onocsinu is evod ,)ocitamsalpirep oizaps(erallulec eterap al e anarbmem al art oizaps ollen acitamsalpotic anarbmem al osrevartta itatropsart e amsalpotic len itazzitetnis onos onaculgoditpep led itnenopmoc isrevid I .pps allenagroM ,.pps retcabortiC ,.pps sanomoduesP ,.pps retcaboretnE oipmese da(acimosomorc enigiro id isamattal-ateb noc imsinagrorcim id osac leN .olozadinortem li emoc ,adiciboreana ocitoibitna ehclauq noc manoertza o enoxairtfec id enoizanibmoc al Ì Ã avitanretla artlaânU .EELB onocudorp ehc imsinagrorcim i iserpmoc ,ortteps oipma ¹Âip led icimttalateb i onos icimenepabrac I.senegotyconom airetsiL e itnetsiser inillicitem ihccocolifats ,ihccocoretne ortnoc evittani onos eniropsolafec el ettuT .S ,suerua .)13oiduts onu ni %86(icitoibitna id eilgimaf ertla da osseps e icittal-ateb i ittut a itnetsiser otilos id onos)anilicasso(anilicitem alla etnetsiser suerua .nim/lm 05 a eroirefni eraluremolg enoizartlif noc itteggos ien icimittal-ateb id esod al eratsuigga oirassecen ¨Ã ,elareneg nI .icitoibitna id ailgimaf atseuq etnemacimihc ecsinifed)3 .onazzilordi ehc icimattal-ateb ied enoiznuf ni isamattal-ateb id ipit itlom onos iC .15»Â94 Ãtilibisnesrepi ad isrevva itteffE.2.)isatsoloc(icitape de)einepotic(ilibisrever icigolotame ibrutsid isracifirev onossoP ed sonugla ne sacituÃ©Ãparet senoicartnecnoc etimrep euq ol ,secev 01 a 3 ed atnemua acilÃ¡fecneotameh arerrab al ed sÃ©Ãvart a nÃ³Âicartenep al ,aegnÃÃnem nÃ³Âicamalfni yah odnauc ,ograbme niS .8002 Â©Ã thgirypoC .52socitÃ³Âibitna ed sailimaf sarto a aicnetsiser al ed setnasuac somsinacem sorto a sodaicosa ri nedeup ,senoisaco ne ,euq setnerefid somsinacem 3 etnaidem etnemacisÃ¡ÃÃb socimÃÃtcalateb sol a aicnetsiser rallorrased nedeup sairetcab saL42aicnetsiser ed somsinaceM.saiboreana sairetcab y sovitisopmarg a etnerf dadivitca ed eceraC .EgI al rop sadaidem nÃ¡Âtse on)sorto y sacirÃÃv senoiccefni ne sanilicineponima ed nÃ³Âicartsinimda al sart ametnaxe ,emrofilibrom ametnaxe(anilicinep ed nÃ³Âicartsinimda al sart setneucerf sÃ¡Ãm saenÃÃÃtuc senoiccaer sal euq atneuc ne renet euq yaH .MBC al a roirefni orep MIC al a roirepus se ocitÃÃÃibitna led nÃ³Âicartnecnoc al odnauc royam ohcum se ocitÃÃibitna otneimatart le etnarud setnetsiser setnatum ed nÃ³Âicceles al ,odal orto roP .E ,silahrratac allexaroM ,eazneulfni sulihpomeaH ,anilicitem al a elbisnes suerua .dadilibisnesrepih noc nanoicaler es on euq sosrevda sotcefE.1sosrevda sotcefE.)menepimi ,amidizatfec ,matcabozat noc anilicarepip(aiparetonom ne ortcepse oilpma ed socimÂ¡Ãtcalateb razilitu nedeup es nÂ©Ãibmat euqnua ,odisÃ³Âculgonima nu y)ocimÂ©Ânepabrac o matcabozat noc anilicarepip ,amipefec(acinÃ³Âmoduespitna y avitisopmargitna dadivitca noc ocimÂ¡Ãtcalateb nu ed nÃ³Âicaicosa al ne etsisnoc radnÂ¡Âtse atuap aL .mutaton muillicineP ognoh led aicneserp al etna aÃÃbihni es ovitluc le arap sodazilitu oyasne ed sobut sol ed sonugla ne suerua succocolyhpatS and otneimicerc le euq Ã³Âvresbo ,8291 and erbmeitpes ne ,neiuq gnimelF rednaxelA a eyubirta es anilicinep al and otneimirbucsed lEnÃ³Âiccudortnl .)soiboreana y sovitisopmarg salricudorp nedeup nÂ©Ãibmat euqnua(sovitagenmarg ne etnemlaicepse ,socimÂ¡tcalateb sol a etnerf aicnetsiser ed omsinacem lapicnirp le atneserpeR .)6 .oidem led (cloxacillin, ceftriaxone, ceftazidima and meropenem)12. Despite this, the impact Ã³clÃnica is not significant, as third generation cefalosporinsÃ³ are normally used for treatment, which also eradicates the carrier status.H. influence and resistance to amoxicillin of about 30%; less than 5% of these strains are resistant to beta-lactamase inhibitors, but are susceptible to third generation cephalosporinsÃ³, carbapenÃ Â® mici and aztreonam.Enteric salmonella: ampicillin resistance of 12%, very low to amoxicillin with clavulÃ acid (1 to 5%) and almost nothing to ceftriaxone39.Shigella sonnei: resistance to ampicillin of 15 to 20% (up to 65% in case of traveler diarrhea)40; amoxicillin resistance with clavulniic acid Ã¨ virtually null41.E. coli in a multi-cÃ Â® entric study that includesÃÂ³ mÃs of 2,000 urocultures from women with uncomplicated cystitis, the rate of resistance to ampicillin Ã¨ was 52%, while the rate of resistance to amoxicillin with clavulÃÃ and cefuroxima¨ was less than 3%42. aeruginosa or from micruginosa Extended spectrum beta-lactamase (BLEE) producing organisms. The beta-lactic ring has a structural similarity to the Ã³ region of the pentapÃ Â® peptide to which these enzymes bind, so Ã¨ can covalently join them and prevent the formation of the cell wall Ã³ (the fig. This type of reaction cannot be predicted with the performance of Ã³ sensitivity tests cutÃÃ.Subjects with penicillin allergy may cross-react with other beta-lactamis, such as cephalosporins (in general about 3-4% of the subjects exposed to cephalosporins Ã¨ allergic and 4-10% of subjects with severe viral allergies to penicillins; the reaction Ã³ crociata with first generation cephalosporinsÃ³ Ã¨ Maggiore) or carbapenÃ Â® mici (3% in ,opmeit led etneidneped etnemetnereferp n³Ã icca noc socit³Ã ibitna sol ed aiporp ,acitsÃÃ retcarac atsE .)dadilibisnesrepih ed emordnÃÃ s y nosnhoJ-snevetS ed emordnÃÃ s ,asotnemacidem erbeif ,ailifonisoe ,serlup anomsodartlifni ,osotnemacidem ojif ametire ,emrofitlum ametire(sodiconocsed somsinacem o)semrofiracitru on sametnaxe(T soticofnil rop adaidem adrater dadiladrater isnes ed n³Ã iccaer ,)oreus led dademrefne acisÃ Ã Ã lc al(sojelpmoconumni ,)laicitsretni sitirfen ,sainepotic ,sisil³Ã³Â mehe(sopreucna rop saddem ratse nedeup y ocamrÂ Ã Ã Ã f led Ã ³Ã icicartsined h27sal ed rited a naterp eS .sisod ertne olavretni led %05 at a 04 roirepus MIC of euq royam T not etneicifus aredisnoc senoiccefni ed aÂROYÃ am al arap ,olpmeje roP .mic al secerc noc arepus ocimÂ Â Would like to see tcalateb ed selautibah sisod ed ed n³Ã icartsinimda al saznacla es q sodijet y ergnas ne ocit³Ã sipled ibay ed n³ÃÃ ibay ed nÃ ³Ã wisaced noc al euicartartartucnecamelc born °Â Ã³ Contact gnin neserated per ohatathein sizing on the name of the name, small noc nosnes sinorcim rop oicni ed sosac sol nE .amatcalb siačni al sernoc socimÂ soil a setsiser nos sapec sal ed %01 nu etnemadamixorpA .etnemavitcepser ,sanames 3 adac atsah y h42 adac³Â icartsinimda us etimeuq ol ,ocit repâ©Â ibitna led adinetsos nâÂ©Ã icarebil of eugisxe c anilicinep al of anilicazneb y an³ÃÂ aaccording to nÂÂÂÂÂÂÂÂÂÂÂ âÂ coicã to a Opelmo inumni ed adidrâÃÊ Ã sallã sacigr©ÂÂ from senoiccaer neneit anilicinep al a aigrela ed airotsih noc sotejus sol ed %01 le olÃÃ s euq atneuc ne renet osicerp sE .oiratinumoc negiro ed sosac sol ne elbanozar yum acituÃ³ÂÂÂÂÂÂ paret© nÃ³ Ã icanu se ocã â â â â â â Â Correct valise c oã c sal noc etrapmoc on euq otartsus nu erbos nÂ©Ã icca us necreje euq ay ,sodarelot neib lareneg ne socamrÃ A Ã f f nosÂ Enhance TcalatebL .)anilicinep al a socigrÂÃ from sotejus %11 le y id alleuq a elimis eterap anu onnah itnetsiser loocla idica irettab I .PBP ellen 'E .2.82manoertza ocimattabonom li e enoizareneg azret id eniropsolafec el erazzilordi id odarg ni onos ehc ,EELB itteddisoc i ,enoizareneg avoun anu osrev imizne itseuq id enoizuloveâl otirovaf ah inneced rep icitoibitna id)osubaâl e(osuâL .)atazzilaiceps acidem epiuqeânu id etrap ad e ocimanidome oiggarotinom otaugeda noc ,otteggos led otamrofni osnesnoc oiverp ,oreiladepso etneibma ni(anillicinep .itnednecsa isod noc enoizazzilibisnesed anu erautteffe id ailgisnoc is ,elibativeni ¨Ã icittal-ateb noc otnemattart li e ovitisop ¨Ã tset li eS .P ad aimeirettab noc itteggos id oppurgottos li audividni iduts ilged onu olos e , Ãtilatrom alled enoizudir al adraugir otnauq rep aiparetonom al arepus non atanibmoc aiparet al ehc ottaf lus onadrocnoc iduts ilg ittuT .acinilc acitarp allen atazzilitu 1Âip al e iciborcimitna id asoremun 1Âip ailgimaf al onocsiutitsoc ,acirettab eralullec eterap alled isetnis alled oidats omitluâlled enoizibiniâl Í Â enoizaâd omsinaccem iuc li ,icimattal-ateb icitoibitna ilG .elanedoud enoizrop amirp alled azzetlaâlla acifirev is ehc ,otnemibrossaâl ecsiunimid e atnellar obic id azneserp aL .)CIM >T(CIM al arepus ocitoibitnaâlled enoizartnecnoc al elauq li etnarud)T(opmet li ¨Ã icimttal-ateb ied acinilc aicaciffeâl noc alerroc oilgem ehc ocimanidocamraf e ocitenicocamraf ecidniãL .itseuq id enoizanibmoc isaislauq o acaidrac aimtira ,enoisnetopi ,oegniral amede ,amedeoigna ,)eenatuc inoisel id ipit irtla non e(otirurp o osnetni erossor ,airacitro emoc isratsefinam 2Âup acigrella enoizaer id opit otseuQ .elamitto CIM alled eroiggam T nu erenetto rep onroig la etlov ¹Âip enoizartsinimmos al airassecen odnedner ,aigolosop al anoizidnoc en ,icimttal-ateb ied etrap roiggam alled enoizanimile id ativime everb al noc microorganismos microorganismos grampositivos, pero con una capa de peptidoglucano fina y, por fuera, una capa muy rica en lÃÃpidos (fig. aeruginosa.3.Tanto los carbapenÃ©Ãmicos como la piperacilina y el tazobactam se reservan para la peritonitis postoperatoria (nosocomial).4.La piperacilina con tazobactam es un antibiÃ³Ático adecuado para la colangitis en sujetos con antecedentes de instrumentaciÃ³Ân o consumo previo de antibiÃ³Âticos, causada habitualmente por enterobacterias, estreptococos, enterococos y, ocasionalmente, por P. En todo caso, la epidemiologÃÃa local debe determinar la pauta empÃÃrica mÃ¡Ãs adecuada, en funciÃ³Ân del patrÃ³Ân de resistencias local.Endocarditis bacteriana1.La penicilina es el antibiÃ³Ático de eleccIÃ³Ân en la endocarditis causada por estreptococos del grupo viridans, y puede sustituirse por la ceftriaxona. La demostrada sinergia de ceftriaxona y ampicilina in vitro se ha aplicado con Ã©Âxito en el tratamiento de la endocarditis enterocÃ³Ãcica23.Mecanismo de acciÃ³Ân10Los antibiÃ³Âticos betalactÃ¡Ãmicos son agentes bactericidas que producen su efecto principalmente a travÃ©Ãs de 2 mecanismos: inhibiciÃ³Ân de la sÃÃntesis de la pared bacteriana e induccIÃ³Ân de la autÃ³Ãlisis bacteriana. catarrhalis). AsÃÃ, de forma somera, los principales mecanismos implicados en la resistencia son los siguientes:1.ProduccIÃ³Ân de enzimas (betalactamasas). Cuanto mÃ¡Ãs elevada sea la CIM, mÃ¡Ãs difÃÃcil serÃ¡Á alcanzar y superar durante el tiempo necesario esa concentraciÃ³Ân tras la administraciÃ³Ân del antibiÃ³Ático. Las bacterias gramnegativas tienen una pared mÃ¡Ãs fina y compleja que consta de una membrana externa formada por lÃÃpidos y proteÃÃnas, y de una capa interna delgada de peptidoglucano. Es por eso que estas enzimas se

llaman tambiÃ©Ân PBP (penicillin binding protein ¢ÃÂproteÃÂna ligada a la penicilina¢ÃÂ). Diferentes alteraciones en las PBP (mutaciones, hiperexpresiÃ³An y modificaciÃ³An de la afinidad) pueden dificultar la uniÃ³An eralocitrap ni(anillicitem alla itnetsiser ihccocolifats ,asonigurea .ailgam anu onamrof e orol art onocsinu is ehc iditpepartet id enetac assif ocimrum odicaâl ,atlov aus A .P ad enoizefni id oihcsir a o ehcigoloru inoizalopinam noc itteggos ieN .PBP ella erinu isretop rep ocitamsalpirep oizaps ollen ossergniâl onitilicaf ehc)irop(enietorp id onatissecen)icimttalateb i emoc(elifopil ocop eznatsos el ,)ivitagen-marg imsinagrorcim ied osac len emoc(eralullec anarbmem anu id azneserp allad atatneserppar areirrab alla etnorf iD .elacol acimedipe enoizautis allad ednepid e , Ãtinu el e irtnec i art etnemloveton airav icitoibitna ilga aznetsiser aL .elirbbef aineportuen noc itteggos ied ociripme otnemattart led elatnemadnof etnenopmoc anu onos icimttalateb Ielirbbef aineportueN.anicimocnav ,etnemlanoisacco ,e edisocilgonima nu da otaicossa)menepimi o matcabozat noc anilicarepip emoc(elanimoduesp-itna Ãtivitta noc ocittal-ateb nu onodulcni adiug eenil elled etrap roiggam aL .S ad ittodorp illeuq onos ipmese inucla ;isamattal-ateb el eravittani da onodnet isamattal-ateb elled irotibini ilg e ilibirefsart eresse onossop esse ,osac omirp leN .enoitseuq ni omsinagrorcim led CIM allad arusim agral ni ednepid ovitteibo otseuq id otnemignuiggar li ehc eripac elicaf ¨Ã ,4 arugif al odnavressO .iboreana ivitagen-marg illicab inucla e etehcorips el ©Ãhcnon ,)iboreana ehc ivitatlocaf ais(ivitisop-marg illicab i e evitagen-marg occoc id icon el ,evitisop-marg occoc id icon el etnemlaizini avednerpmoc G anillicinep alled ociborcimitna ortteps oL.eterap alled ehcitsiretarac ella atuvod atnel enoizartenep anu da atinu etnemlibaborp ,isamattal-ateb id enoizudorp alla atuvod Ì Ã irettabocim ied aznetsiser elarutan aL .isac ied %57 len ecaciffe ¨Ã de elanoizecce ¨Ã issalifanaâl am ,)itteggos ied ozret nu da onif(icigrella inemonef eracifirev onossop is arudecorp atseuq noC . Ãtivitta aus al ecsiunimid ehc ,enietorp ella ocittalateb id id azneserp catheters) and other multi-resistant bacteria, according local . They have a certain irritant action at the site of administration and may cause gastritis if administered digestively, pain with intramuscular injection (type of injectable bencilpenicillin preparations) or phlebitis (characteristic of some betalactamics, such as cloxa (ciline). After sensitization, drug causes much faster clinical manifestations (minutes to hours). Without the wall, the bacterium is exposed to the medium and dies due to changes in the birth pressure. Although there have been no truly new beta-lactic drugs available for more than two decades¹, relentless increase in resistance and advancement in knowledge of their molecular mechanisms have conditioned existence of a large amount information in the medical literature on each of the components of this antibiotic family. Three of those. In some preparations, such as cefoperazone or ceftriaxone, bile predominates. In cases (confirmed or suspected) of an IgE-mediated reaction, when it is necessary to administer a betalactic (or earlier if possible), Ã¨ it is advisable to perform a skin sensitivity test to confirm the diagnosis. Betalactymics have wide body , with appropriate serum and tissue concentrations in most tissues, including bile and synovial fluid; they cross the placental barrier without problems, but do not penetrate well into the non-inflamed central nervous system and eyes. Events allergic to penicillin can be classified, depending on mechanism and chronology, into:(a) Immunoglobulin E IgE)-mediated reactions, which are generally immediate (usually within the first 15 minutes), but up to 5% ed serotcudorp somsynagroorcim ed aicneucerf royam(acinÃ³Ã rc avitcurtsbo ranomlup dademrefne noc sotejus aÃ nomuen ed sosac sol y)oiratinas ametsis la adaicosa aÃ nomuen ed sosac sol ed sohcum a Ã"Ã Ra lcÃ"Ã ogseir ed serotcaf sol noc sotejus a otnat aÃ riulni euq(n³Ã wing rop a a ÂÃ nomuen ed sosac sol elbireferenp Age luvalc odicÃ Â Â Â Advise noc anilicixoma noc noc otneimatart lE .rojem se)selaro saniropsolafec o anilicixoma al omoc(socit©Ã³Ã ticinisimes y socitÃ©Ã tnis soired davired at ed nÃ ³ÃÃ icrosba al q poor tim)ird aced sal³adged asifAEDAÁ rollÃ Â Roll it Â Â Â Â Â Â They aditid a brosba saicnatsus saS to l .I rop al aiaion igr© Increase to senoiccaer sal ne eyulfni on y ocitÃ³Ã ibitna led osu areiuqer es euq zev adac olriteper osicerp sE .anilical a elbissuerua .retcaboretn e osac le acim³Ã³Ã somorc asamatcalb ed nÃ ³Ã eraped of a osac remirmirp le acidÃ a mã salp ed nÃ ³E desed of the idebed nÃ³ netsiser ririuqda nedeup y sanilicinep sal a aicnetsiser larutan amrof ed natneserp:.pps retcaboretnE y eainomuenp .K.44,34)%5,5 la 7,1(EELB ed arotcudorp se sapec ed royam zev adac nÃ³This ropU .anoloniuq anu a odiâÂ44,34)%5,5 la 7,1(EELB ed arotcudorp se sapec ed royam adac nÃÃ iciced the ocicom Ã Ã Ã Ã Tcalatele oiralatipsosososernei qer euq ogseir otla ed sosac sol nE .S(somsinagroorcim sonugla ed erop sasamatcalateb ed nÃ³ÃÃÃ cudorp al se socimÂ Ã Ã tcalateen b etneretnerf natropmi sÃ Ã Ã Ã Ã